

Marked Length Rigidity For One Dimensional Spaces

Jean-François Lafont

February 1, 2008

Abstract

In this paper, we give a proof of the marked length spectrum (MLS) rigidity for complete geodesic metric spaces of topological dimension one. More precisely, in a compact geodesic metric space of topological dimension one, the minimal length of a loop in a free homotopy class is well defined, and provides a function $l : \pi_1(X) \longrightarrow \mathbb{R}^+ \cup \infty$ (the value ∞ being assigned to loops which are not freely homotopic to any rectifiable loops).

We introduce a subset $Conv(X)$, which is merely the union of all minimal loops of finite length. Now let X_1, X_2 be a pair of complete, compact, geodesic metric spaces of topological dimension one, and define $Y_i = \overline{Conv(X_i)}$. Then we show that any isomorphism $\phi : \pi_1(X_1) \longrightarrow \pi_1(X_2)$ satisfying $l_2 \circ \phi = l_1$, forces the existence of an isometry $\Phi : Y_1 \longrightarrow Y_2$ which induces the map ϕ when restricted to the fundamental groups of the Y_i .

1 Introduction.

This paper is motivated by a long standing conjecture concerning negatively curved manifolds: that the length of closed geodesics on a negatively curved Riemannian manifold determines the space up to isometry. More precisely, in a negatively curved manifold (M^n, g_i) , there are unique geodesics in free homotopy classes of loops, and we can consider the length function $l : \pi_1(M^n) \longrightarrow \mathbb{R}^+$. The marked length spectrum (MLS) conjecture states that, if we have a pair of negatively curved Riemannian metrics on M^n which yield the same length function l , then they are in fact isometric. The conjecture has been proved for certain specific manifolds, most notably for 2-dimensional manifolds (independantly by Croke [5] and Otal [12]) as well as for rank one locally symmetric spaces (by Hamenstadt [10]), and higher rank locally symmetric spaces (by Dal'bo & Kim [9]).

In this paper, we consider geodesic spaces of topological dimension one. The starting observation is that these spaces share a lot of the properties of negatively curved manifolds. In particular, they are aspherical (see Curtis & Fort [8]), and have unique minimal length representatives in each free homotopy class of loops (by Curtis & Fort [7], also shown by Cannon & Conner [4]). In particular, one can ask whether the MLS conjecture holds in the setting of geodesic spaces of topological dimension one. We define a subset $Conv(X)$ of any geodesic space X of topological dimension one, and show the following theorem:

Theorem 1.1. *Let X_1, X_2 be a pair of complete, compact, geodesic spaces of topological dimension one, and define $Y_i = \overline{Conv(X_i)}$. Assume the two spaces have the same marked length spectrum, that is to say, there exists an isomorphism $\phi : \pi_1(X_1) \longrightarrow \pi_1(X_2)$ satisfying $l_2 \circ \phi = l_1$. Then Y_1 is isometric to Y_2 , and the isometry induces (up to change of basepoints) the isomorphism ϕ on the subgroups $\pi_1(Y_i)$.*

Let us provide an outline of the proof, with reference to the next section for appropriate definitions. The idea behind the argument is to look at a certain subset of $Conv(X_1)$ consisting of *branch points*. For a pair of branch points, one can look at a minimal geodesic joining them, and one constructs a pair of geodesic loops with the property that they intersect precisely in the given minimal geodesic. Now using the isomorphism of fundamental groups, one can look at the corresponding pair of geodesic loops in $Conv(X_2)$. Using the fact that the lengths are preserved, one shows that the corresponding pair in $Conv(X_2)$ likewise intersects in a geodesic segment, and that furthermore, the length of the intersection is exactly equal to the length of the original geodesic segment. One can then proceed to show that this correspondance is well-defined (i.e. does not depend on the pair of geodesic loops one constructs), and preserves concatenations of geodesic segments. This provides us with an isometry between the sets of branch points. Using completeness, we can

extend this to an isometry between the closures of the sets of branch points. Finally, we can consider points in $\text{Conv}(X_1)$ which are *not* in the closure of the set of branch points, and it is easy to see that each of these points lies on a unique geodesic segment with the property that the segment contains precisely two branch points, which are its endpoints. The correspondance between geodesic segments shows that there is a unique, well-defined, corresponding segment in $\text{Conv}(X_2)$ of precisely the same length, allowing us to extend our isometry to all of $\text{Conv}(X_1)$. One then extends the isometry to the closure, obtaining that the two spaces Y_i are isometric. Finally, it is clear from the construction that the isometry induces the isomorphism between the fundamental groups of the Y_i (the fact that $\pi_1(Y_i)$ injects in $\pi_1(X_1)$ follows from a result of Curtis & Fort [7]).

Acknowledgments.

The results contained in this paper were part of the author's thesis, completed at the Univ. of Michigan, under the guidance of professor R. Spatzier. The author would like to thank his advisor for his help throughout the author's graduate years. The author would also like to thank professors H. Bass and J. Heinonen for taking the time to proofread his thesis and to comment on the results contained therein.

2 Preliminaries.

In this section, we will define the various terminology used in this paper, as well as quote certain basic results we will use in our proofs. We start by reminding the reader of certain basic notions on length spaces (and refer to Burago, Burago & Ivanov [3] for more details on the theory).

Definition 2.1. A *path* in a metric space (X, d) , is a continuous map $f : [a, b] \longrightarrow X$ from a closed interval into X . A *loop* in X is a path f satisfying $f(a) = f(b)$. Observe that we can always view a loop as a based continuous map from $(S^1, *)$ to $(X, f(a))$.

Definition 2.2. Let (X, d) be a metric space. The *induced length structure* is a function on the set of paths, denoted by l_d , and defined as follows:

$$l_d(\gamma) := \sup \sum_{i=1}^n d(\gamma(x_{i-1}), \gamma(x_i))$$

where $\gamma : [a, b] \longrightarrow X$ is a path, and the supremum ranges over all finite collection of points $a = x_0 < x_1 < \dots < x_n = b$. We will say a path γ is *rectifiable* provided $l_d(\gamma) < \infty$.

Observe that rectifiability is preserved under finite concatenation of paths, and under restriction to subpaths.

Definition 2.3. Let (X, d) be a metric space, and l_d the induced length structure. We define the *intrinsic pseudo-metric* $\hat{d}$ induced by d as follows. Let $p, q \in X$ be an arbitrary pair of points, and define the $\hat{d}$ distance between them to be $\hat{d}(p, q) := \inf l_d(\gamma)$, where the infimum ranges over all paths $\gamma : [a, b] \rightarrow X$ satisfying $\gamma(a) = p$, $\gamma(b) = q$.

Note that the function $\hat{d}$ actually maps $X \times X$ to $[0, \infty]$, where two points $p, q \in X$ have $\hat{d}(p, q) = \infty$ if and only if there are no rectifiable paths joining p to q .

Definition 2.4. Let (X, d) be a metric space, $\hat{d}$ the corresponding intrinsic pseudo-metric. We call (X, d) a *length space* if $d = \hat{d}$ (in particular, $\hat{d}$ has image in $(0, \infty)$).

Definition 2.5. Let (X, d) be a length space. We say that (X, d) is a *geodesic space* if, for every pair of points $p, q \in X$, there is a path $\gamma_{p,q}$ joining p to q , and having length precisely $d(p, q)$. Such a path is called a *distance minimizer*.

In topology, one of the most important concepts is that of dimension. While there are many different notions of dimension, the one which will be of interest to us is that of *covering dimension*.

Definition 2.6. Let X be a topological space. We say that X has covering dimension $\leq n$ if, for any open covering $\{U_i\}$ of X , there is a refinement $\{V_i\}$ with the property that every $x \in X$ lies in at most $n + 1$ of the V_i . We say that X is *n-dimensional* if X has covering dimension $\leq n$, but does not have covering dimension $\leq n - 1$.

We will denote the covering dimension of a space X by $Cdim(X)$. Observe that any path connected topological space has $Cdim(X) \geq 1$. The spaces we will be interested in are those satisfying $Cdim(X) = 1$. Examples of such spaces are plentiful. In particular, we have the following criterion (see Chapter VII in Hurewicz & Wallman [11]):

Theorem 2.1. *Let (X, d) be a metric space, and denote by $Hdim(X)$ the Hausdorff dimension of X . Consider the induced topology on X . One has the inequality $Cdim(X) \leq Hdim(X)$.*

Hence any path connected metric space with Hausdorff dimension less than two has covering dimension one. We will henceforth, unless explicitly mentioned otherwise, focus exclusively on geodesic spaces of topological dimension one. Note that,

in general, a one dimensional geodesic space might not be non-positively curved ¹. However, such spaces often exhibit properties which are quite similar to those of negatively curved spaces. In order to show the property we are interested in, we start by providing a few definitions.

Definition 2.7. Let $\gamma : (S^1, *) \longrightarrow (X, p)$ be a loop. We say that the loop is *reducible* provided that there is an open interval $I = (x, y) \subset S^1 - \{*\}$ such that $\gamma(x) = \gamma(y)$ and the loop $\gamma|_{[x, y]}$ is nullhomotopic. We say that γ is *cyclically reduced* if the interval I is allowed to include the basepoint $*$. A curve which is not reducible (resp. cyclically reducible) will be said to be *reduced* (resp. *cyclically reduced*). We define a constant loop to be cyclically reduced.

Definition 2.8. Let (X, d) be a 1-dimensional geodesic space. We say that a loop γ is a *geodesic loop* provided γ is rectifiable and is the unique length minimizer in its free homotopy class. We say that a path is *geodesic* provided that it is reduced and rectifiable.

WARNING: Our terminology differs from the standard, in that our geodesics might not even be locally minimizing.

Note that the definition of a reduced loop applies equally well to a path, so that we can talk about reduced paths in a geodesic space. We now state Cannon & Conner's theorem (Thm 3.9 in [4]):

Theorem 2.2 (Uniqueness of geodesic loops). *Let (X, d) be a compact, path-connected, 1-dimensional metric space. Then*

- *every loop is homotopic to a reduced loop, which is unique up to reparametrization,*
- *every loop is freely homotopic to a cyclically reduced loop, which is unique up to cyclic reparametrization.*

Using this result, it is easy to show that in every free homotopy class of loops that contains a rectifiable curve, there is a unique minimal length representative. Note that any length space is automatically a path connected metric space.

Corollary 2.1 (Minimality of geodesic loops). *Let (X, d) be a compact 1-dimensional length space, γ a loop. Then one of the following two possibilities holds:*

¹Recall that negatively curved metric spaces are defined by the property that their universal cover satisfies the $CAT(-\delta^2)$ condition (a metric version of negative curvature, see Bridson & Haefliger [2]). Naturally, the problem in our setting is that spaces which are very singular might not have a universal cover.

- *there is no rectifiable loop freely homotopic to γ ,*
- *there is a unique (up to reparametrization) minimal length rectifiable loop freely homotopic to γ .*

Proof:. Let us assume that there is a rectifiable loop freely homotopic to γ , denote it by $\bar{\gamma}$. Assume that $\bar{\gamma}$ is not cyclically reduced. Then there exists an interval $I = [x, y] \subset S^1$ with $\bar{\gamma}(x) = \bar{\gamma}(y)$, such that the loop $\bar{\gamma}_1$ obtained by restricting $\bar{\gamma}$ to $[x, y]$ is contractible. Consider the loop $\bar{\gamma}_2$ defined by restricting $\bar{\gamma}$ to the closure of $S^1 - I$. Then by contracting $\bar{\gamma}_1$, we see that $\bar{\gamma}$ is freely homotopic to $\bar{\gamma}_2$. Furthermore, we have that $l(\bar{\gamma}) = l(\bar{\gamma}_1) + l(\bar{\gamma}_2) > l(\bar{\gamma}_2)$. Hence $\bar{\gamma}$ cannot be of minimal length.

This forces any minimal length loop in a free homotopy class to be cyclically reduced. But by the theorem of Cannon & Conner, such a loop is unique up to reparametrization. Our claim follows.

It is now straightforward to prove an analogue of the Cannon & Conner result for paths in a 1-dimensional metric space. This statement is our following:

Proposition 2.1 (Uniqueness of geodesic paths). *Let (X, d) be a compact, path-connected, 1-dimensional metric space. Then every path is homotopic (relative to the endpoints of the path) to a reduced path, which is unique up to reparametrization.*

Proof:. Let X be our geodesic space, $\mathbf{p}$ our path, and p, q the two endpoints. Consider the space X' , defined to be the quotient space of X obtained by identifying p and q . We claim that X' is a 1-dimensional geodesic space. Indeed, quotienting can only decrease the dimension of a space, hence $Cdim(X') \leq 1$, and quotients of path connected spaces are path connected, so $Cdim(X') \geq 1$, resulting in X' being 1-dimensional. Furthermore, X' is a compact, path-connected metric space, with the image $\mathbf{p}'$ of $\mathbf{p}$ a closed loop based at the point $p = q$. By the first part of Theorem 2.2, we have that this loop is homotopic to a unique (upto reparametrization) reduced loop. As there is a bijective correspondance between homotopies of $\mathbf{p}'$ which preserve the basepoint and endpoint preserving homotopies of $\mathbf{p}$, we immediately get our claim.

Corollary 2.2 (Minimality of geodesic paths). *Let (X, d) be a compact 1-dimensional length space, $\mathbf{p}$ a rectifiable path joining points p and q . Then the unique reduced path joining p to q homotopic (relative to the endpoints) to $\mathbf{p}$ has minimal length amongst all paths with this property.*

Proof:. This argument is identical to the one for loops: assume that $\mathbf{p}$ is rectifiable, but not reduced, and is a mapping from $[a, b]$ into X . Then there is a subinterval $I = [a', b'] \subset [a, b]$ such that $\mathbf{p}(a') = \mathbf{p}(b')$ and the loop $\bar{\mathbf{p}}_1$ obtained by restricting $\bar{\mathbf{p}}$

to I is contractible. Denoting by $\mathbf{p}_2$ the concatenation of $\mathbf{p}$ restricted to $[a, a']$ and $[b', b]$, we see that $\mathbf{p}_2$ is homotopic to $\mathbf{p}$ via a homotopy preserving the endpoints. Furthermore, $l_d(\mathbf{p}_2) < l_d(\mathbf{p})$, which yields our claim.

Corollary 2.2 immediately yields:

Corollary 2.3. *Let (X, d) be a compact 1-dimensional geodesic space, γ a distance minimizer joining p to q . Then γ is a geodesic.*

Definition 2.9. Given a geodesic space X of topological dimension one, we define the π_1 -hull of X , denoted $\overline{Conv(X)}$, as the union of (the images of) all geodesic loops in X . A space satisfying $X = \overline{Conv(X)}$ is said to be π_1 -convex.

We fix the following notation: given a pair of paths $\mathbf{p}$ and $\mathbf{q}$, with the terminal endpoint of $\mathbf{q}$ coinciding with the initial endpoint of $\mathbf{p}$, we will denote by $\mathbf{p} * \mathbf{q}$ the concatenation of the two paths (traversing $\mathbf{q}$ first, followed by $\mathbf{p}$), and by $\mathbf{p}^{-1}$ the path obtained by reversing $\mathbf{p}$. We start by proving several lemmas concerning concatenations of various types of paths. Throughout the rest of this section, we will assume that the underlying space X we are working in is a compact, 1-dimensional length space.

Lemma 2.1 (Concatenation of reduced paths). *Let $\mathbf{p}_1, \mathbf{p}_2$ be a pair of reduced paths, parametrized by arclength, in X . Let $t_i = l(\mathbf{p}_i)$ be the length of the respective paths, and assume that $\mathbf{p}_1(t_1) = \mathbf{p}_2(0)$ (i.e. that they have a common endpoint). Then any reduced path $\mathbf{q}$ homotopic to $\mathbf{p} := \mathbf{p}_2 * \mathbf{p}_1$ is of the form $\mathbf{q}_2 * \mathbf{q}_1$ where $\mathbf{q}_i$ is a subpath of $\mathbf{p}_i$. Furthermore, we have a decomposition $\mathbf{p}_1 = \mathbf{r}_1 * \mathbf{q}_1$ and $\mathbf{p}_2 = \mathbf{q}_2 * \mathbf{r}_2$, satisfying $\mathbf{r}_1 = \mathbf{r}_2^{-1}$.*

Proof. Let us define $\mathbf{p}$ to be the concatenation $\mathbf{p}_2 * \mathbf{p}_1$. We start by observing that the claim is trivial if the concatenation $\mathbf{p}$ is reduced (just take $\mathbf{q}_i = \mathbf{p}_i$). So let us assume that the concatenation $\mathbf{p}$ is *not* a reduced path, and view it as a map from $D := [0, t_1 + t_2]$ into X . Since this path is reducible, there exists closed intervals $U_j \subset D$ with the property that $\mathbf{p}$ restricted to each U_j is a closed path which is null homotopic. Since each of the paths $\mathbf{p}_1, \mathbf{p}_2$ is reduced, this forces $t_1 \in U_j$.

We now claim that, under inclusion, the family of such closed intervals forms a totally ordered set. In order to see this, we show that any such set $U_j = [a_j, b_j]$ is a symmetric closed interval around t_1 (i.e. that $(a_j + b_j)/2 = t_1$). But this is clear: one can just consider the restriction of $\mathbf{p}$ to the two sets $[a_j, t_1]$ and $[t_1, b_j]$. This yields a pair of paths, parametrized by arclength, joining the point $\mathbf{p}(a_j) = \mathbf{p}(b_j)$ to the point $\mathbf{p}(t_1)$. Furthermore, each of these paths is reduced (since they are subpaths of the reduced paths $\mathbf{p}_1$ and $\mathbf{p}_2$ respectively). But we know that there is a unique reduced path in each endpoint preserving homotopy class of paths joining a pair of points.

Hence the two paths have to coincide, and as they are parametrized by arclength, we immediately obtain our claim.

Next, we claim that this totally ordered chain has a maximal element. Indeed, consider the set U defined to be the union of our sets U_j . We claim that U is still within our family. To see this, we merely note that, by our previous observation on the $U_j = [a_j, b_j]$, the restriction of $\mathbf{p}$ to each symmetric (about t_1) subinterval of U consists merely of traversing some reduced path on $[a_j, t_1]$, followed by backtracking along the same path on $[t_1, b_j]$. By continuity, the same must hold for the symmetric closed interval $\bar{U}$, so that the closure of U also lies within our family. Hence $U = \bar{U}$, and we have found our maximal element.

It is now easy to complete our proof: if $[a, b]$ is our maximal interval U , we can now define our $\mathbf{q}_1$ and $\mathbf{r}_1$ explicitly. We set $\mathbf{q}_1 := \mathbf{p}|_{[0, a]}$, $\mathbf{r}_1 := \mathbf{p}|_{[a, t_1]}$, $\mathbf{r}_2 := \mathbf{p}|_{[t_1, b]}$, and $\mathbf{q}_2 := \mathbf{p}|_{[b, t_1 + t_2]}$. We note that it is clear that $\mathbf{p}_1 = \mathbf{r}_1 * \mathbf{q}_1$ and $\mathbf{p}_2 = \mathbf{q}_2 * \mathbf{r}_2$. From our proof, it is also immediate that $\mathbf{r}_1 = \mathbf{r}_2^{-1}$. Finally, since U was picked to be maximal, the path $\mathbf{q}_2 * \mathbf{q}_1$ must be reduced. Which was precisely our original claim.

Corollary 2.4 (Form of reduced loops). *Let η be a reduced loop in X . Then η can be expressed as a concatenation $\mathbf{p}^{-1} * \gamma * \mathbf{p}$, where $\mathbf{p}$ is a reduced path, and γ is a geodesic loop.*

Proof. Let us view η , parametrized by arclength, as a map from $[0, t]$ into X . Consider the point $p := \eta(t/2)$, and consider the pair of paths $\mathbf{q}_1 := \eta|_{[0, t/2]}$ and $\mathbf{q}_2 := \eta|_{[t/2, t]}$. Observe that each of these paths is reduced (being a subpath of η), have common endpoints, and that $\eta = \mathbf{q}_2 * \mathbf{q}_1$. Now consider the concatenation of paths $\mathbf{q}_1 * \mathbf{q}_2$ and apply the previous theorem. Our claim immediately follows.

Lemma 2.2 (Path-loop-path concatenations). *Let $\mathbf{p}$ be a reduced path parametrized by arclength $[0, t]$, γ a cyclically reduced loop based at $p := \mathbf{p}(t)$, parametrized by arclength $[0, s]$. Then the unique reduced loop η in the homotopy class of $\mathbf{p}^{-1} * \gamma * \mathbf{p}$ must pass through the point p . Furthermore, this loop either coincides with $\mathbf{p}$ on the interval $[0, t]$, or coincides with $\mathbf{p}^{-1}$ on the interval $[l(\eta) - t, l(\eta)]$.*

Proof. Assume not, and let us start by focusing on $\gamma * \mathbf{p}$. Since $(\gamma * \mathbf{p})(t) = p$, our reduced loop η must avoid it, which forces $\gamma * \mathbf{p}$ to be reducible. By the previous lemma, this implies that γ^{-1} restricted to $[s - \epsilon, s]$ must coincide exactly with $\mathbf{p}$ restricted to $[t - \epsilon, t]$. But by definition of γ^{-1} , we see that the restriction of γ^{-1} to $[s - \epsilon', s]$ coincides with γ restricted to $[0, \epsilon']$. Applying the same analysis to $\mathbf{p}^{-1} * \gamma$ gives that γ restricted to some $[s - \epsilon', s]$ must coincide with $\mathbf{p}$ restricted to $[t - \epsilon', t]$.

Picking δ smaller than both ϵ and ϵ' , we get that γ restricted to $[0, \delta]$ coincides with γ restricted to $[s - \delta, s]$. But this contradicts the assumption that γ was cyclically reduced. The first claim follows.

For the second claim, we observe that the argument we just gave in fact shows that the intervals $U \subset [0, 2t + s]$ that give a reduction of the concatenated loop $\mathbf{p}^{-1} * \gamma * \mathbf{p}$ either contain the point t , or contain the point $t + s$, but not both. In particular, this forces η to either start by travelling along $\mathbf{p}$, or to finish by travelling along $\mathbf{p}^{-1}$, which is precisely the second claim.

Note that, in our previous lemma, we can always (by reversing γ if need be) get the reduced loop η to coincide with $\mathbf{p}$ on the interval $[0, t]$.

Lemma 2.3 (Extension of geodesic paths to geodesic loops). *Every geodesic path in X whose endpoints lie in $\text{Conv}(X)$ can be extended to a geodesic loop.*

Proof. Let $\mathbf{p}$ be a geodesic path, parametrized by arclength $[0, t]$. Let us subdivide our path at the point $\mathbf{p}(t/2)$, to obtain a concatenation $\mathbf{p} = \mathbf{p}_2 * \mathbf{p}_1$. Now we can consider the pair of endpoints $\mathbf{p}(0)$ and $\mathbf{p}(t)$, and note that since our geodesic lies in $\text{Conv}(X)$, there are a pair γ_1 and γ_2 of geodesic loops passing through $\mathbf{p}(0)$ and $\mathbf{p}(t)$ respectively.

We can now consider the concatenations: $\mathbf{p}_1 * \gamma_1 * \mathbf{p}_1^{-1}$ and $\mathbf{p}_2^{-1} * \gamma_2 * \mathbf{p}_2$. Consider the unique reduced loops η_1 and η_2 in the corresponding homotopy class. By our previous lemma, they must pass through $\mathbf{p}(0)$ and $\mathbf{p}(t)$ respectively. Furthermore, by reversing the reduced loops if needed, we have that $\eta_1^{-1}|_{[l(\eta_1)-t/2, l(\eta_1)]} = \mathbf{p}_1|_{[0, t/2]} = \mathbf{p}|_{[0, t/2]}$, while $\eta_2|_{[0, t/2]} = \mathbf{p}_2|_{[0, t/2]} = \mathbf{p}|_{[t/2, t]}$. Concatenating these, we obtain a closed curve $\eta_1^{-1} * \eta_2$ which might not be geodesic, but certainly passes through $\mathbf{p}$. Now consider the geodesic loop in the corresponding free homotopy class. We claim that it must still pass through $\mathbf{p}$. Indeed, if it did not, there would be a reduction along some open set containing the point $\eta_1^{-1}(l(\eta_1)) = \eta_2(0) = \mathbf{p}(t/2)$. But as we have seen, any such reduction would have to occur along a symmetric interval around the point, corresponding to some subpath $\mathbf{r}^{-1} * \mathbf{r}$ within $\mathbf{p}$, contradicting the fact that the latter is reduced. Our claim follows.

To illustrate the usefulness of the previous lemmas, we note the following immediate corollary:

Corollary 2.5. *Let X be a geodesic space of topological dimension one. Then $\text{Conv}(X)$ is path connected. Furthermore, $\text{Conv}(X)$ is a strongly convex subset of X .²*

Proof. Let $p, q \in \text{Conv}(X)$ be an arbitrary pair of distinct points, and let $\mathbf{p}$ be a distance minimizer joining the two points. Since $\mathbf{p}$ is a distance minimizer, it is geodesic. So by the previous lemma, there is a geodesic loop extending it. This immediately shows that $\mathbf{p}$ itself lies in $\text{Conv}(X)$. Both our claims follow.

²We say that a subset of a geodesic space is strongly convex provided that for every pair of points in the subset, every distance minimizer joining them also lies within the subset.

Note that in any free homotopy class of loops, we can consider the unique cyclically reduced loop. Then we either have:

- the loop is rectifiable and is a geodesic loop, or
- no freely homotopic loop is rectifiable.

As such, when looking for geodesic loops, it is sufficient to restrict throughout to cyclically reduced loops. We finish this section by defining the marked length spectrum.

Definition 2.10. Let (X, d) be a geodesic space. Assume that, in each free homotopy class of curves on X , there is at most one minimal length representative. The *marked length spectrum* (abbreviated to MLS) is defined to be the function $l_d : \pi_1(X) \longrightarrow \mathbb{R}^+ \cup \infty$ which assigns to each element of $\pi_1(X)$ the length of the corresponding minimal length loop (and assigns ∞ to the free homotopy classes that contain no rectifiable representatives).

Observe that, for geodesic spaces of topological dimension one, by Theorem 2.1 we have that the marked length spectrum is defined.

3 MLS Rigidity for Spaces of Topological Dimension One

For the convenience of the reader, we restate our main theorem:

Theorem 3.1. *Let X_1, X_2 be a pair of complete, compact, geodesic spaces of topological dimension one, and define $Y_i = \overline{\text{Conv}(X_i)}$. Assume the two spaces have the same marked length spectrum, that is to say, there exists an isomorphism $\phi : \pi_1(X_1) \longrightarrow \pi_1(X_2)$ satisfying $l_2 \circ \phi = l_1$. Then Y_1 is isometric to Y_2 , and the isometry induces (up to change of basepoints) the isomorphism ϕ on the subgroups $\pi_1(Y_i)$.*

Throughout this section, we will restrict our attention to spaces X which are complete, compact, geodesic, of topological dimension one, and we set $Y = \overline{\text{Conv}(X)}$. We start by introducing a few definitions and proving an important lemma.

Definition 3.1. Let X be a 1-dimensional geodesic space, p a point in X . We say that X has *branching* at p provided there exists a triple of geodesic paths $\gamma_i : [0, \epsilon] \longrightarrow X$ with the following properties:

- $\gamma_i(0) = p$ for all three paths,

- each concatenated path $\gamma_i * \gamma_j^{-1}$ is a geodesic path.

Note in particular that each of the concatenations must be a reduced path. If X has branching at p , we call p a *branch point* of X .

Definition 3.2. Given a geodesic path $\mathbf{p}$ in a 1-dimensional geodesic space X joining branch points p to q , we say that a pair of geodesic loops γ_1, γ_2 based at p and parametrized by arc-length are $\mathbf{p}$ -distinguishing provided that the two geodesic loops agree on the interval $[0, l(\mathbf{p})]$ (where $l(\mathbf{p})$ is the length of the path $\mathbf{p}$), but on no larger interval. Furthermore, we require that $\gamma_i|_{[0, l(\mathbf{p})]} = \mathbf{p}$ (this condition just means that the geodesic loops start out by respecting the orientation on $\mathbf{p}$).

The importance of $\mathbf{p}$ distinguishing loops lies in the fact that, if γ_1 and γ_2 are $\mathbf{p}$ distinguishing, and if we use an overline to denote the geodesic loop freely homotopic to a given loop, then we automatically have:

$$l(\overline{\gamma_2 * \gamma_1^{-1}}) = l(\gamma_1) + l(\gamma_2) - 2l(\mathbf{p}).$$

In particular, since the concatenated loop represents the product of the elements corresponding to γ_i in $\pi_1(X, \mathbf{p}(0))$, we see that the length of the geodesic path $\mathbf{p}$ can be recovered from the marked length spectrum. We now show the existence of distinguishing geodesics for paths whose endpoints are branching.

Lemma 3.1 (Existence of distinguishing loops). *Let $\mathbf{p}$ be a geodesic path joining a pair of branch points lying in $\text{Conv}(X)$. Then there exists a pair of $\mathbf{p}$ -distinguishing geodesic loops.*

Proof: Let $t := l(\mathbf{p})$. Since our path $\mathbf{p}$ joins a pair of branch points, it is easy to see that there are a pair of geodesic paths γ_1, γ_2 in X which intersect precisely in $\mathbf{p}$. Indeed, let us start by considering $p = \mathbf{p}(0)$, and note that since this point is branching, there exist a triple of geodesic paths $\gamma_i : [0, \epsilon] \rightarrow X$ emanating from the point p with the property that each concatenated path $\gamma_i * \gamma_j^{-1}$ is a geodesic path. Now consider the three possible concatenations $\mathbf{p} * \gamma_i^{-1}$. We claim that at least two of them have to be geodesic paths. Indeed, if not, then two of these concatenations, say $\mathbf{p} * \gamma_1^{-1}$ and $\mathbf{p} * \gamma_2^{-1}$ have to be reducible. Using our previous lemma, this forces $\mathbf{p}$, γ_1 , and γ_2 to coincide in a small enough interval $[0, \epsilon]$. But this contradicts the fact that $\gamma_2 * \gamma_1^{-1}$ is reduced. So we can extend $\mathbf{p}$ past p in two distinct ways, and still have a reduced path. Similarly, we can extend $\mathbf{p}$ past the point $\mathbf{p}(t)$ in two distinct ways, and still have a reduced path. This gives us a pair of geodesic segments which are distinct, then come together and agree precisely along $\mathbf{p}$, and then separate again.

Shrinking the two geodesic segments if need be, we can assume that they are defined on $[-\epsilon, t + \epsilon]$, and that the geodesic $\mathbf{p}$ corresponds to the image of $[0, t]$ in




Figure 1: Initial segments forced to agree: minimal representative of composite curve in first picture has shorter length than corresponding one in the second picture.

both geodesics. We want to extend these geodesic paths to closed geodesic loops. But that is precisely what Lemma 2.3 guarantees. Hence we have a pair of $\mathbf{p}$ distinguishing segments, and we are done.

Before proving our main proposition, we give one last definition.

Definition 3.3. Let X be a compact geodesic space of topological dimension one, and $\mathbf{p}_1, \mathbf{p}_2$ a pair of geodesics in the space, parametrized by the intervals $[a_1, b_1]$ and $[a_2, b_2]$ respectively. We say that $\mathbf{p}_1$ and $\mathbf{p}_2$ are *incident*, provided that $\mathbf{p}_1(b_1) = \mathbf{p}_2(a_2)$. We say that they are *geodesically incident* provided that, in addition to being incident, the concatenated path $\mathbf{p}_2 * \mathbf{p}_1$ is geodesic.

Proposition 3.1. Let X_1, X_2 be a pair of π_1 -convex spaces. If they have the same marked length spectrum, then there is an isometry between the set of branch points of $\text{Conv}(X_i)$.

Proof. We begin by defining a bijective map from $P(X_1)$ to $P(X_2)$, where $P(X_i)$ is the set of geodesic paths joining branch points of $\text{Conv}(X_i)$. Let $\mathbf{p} \in P(X_1)$ be given. By Lemma 3.1, there exists a pair of $\mathbf{p}$ -distinguishing geodesic loops; call them γ_1 and γ_2 . Without loss of generality, we can assume the base point p_1 for $\pi_1(X_1, p_1)$ is the common vertex $\gamma_i(0)$. Now corresponding to the homomorphism $\Phi : \pi_1(X_1, p_1) \longrightarrow \pi_1(X_2, p_2)$, we can find a pair of closed geodesic paths $\Phi(\gamma_1)$ and $\Phi(\gamma_2)$ (i.e. reduced paths, but not necessarily cyclically reduced) based at $p_2 \in G_2$ having precisely the same lengths of minimal representatives in their free homotopy class. We will use an overline to denote the geodesic loop (i.e. cyclically reduced loop) in the free homotopy class of a curve. Observe that by our choice of γ_1, γ_2 being $\mathbf{p}$ -distinguishing, we have that:

$$l_1(\overline{\gamma_2 * \gamma_1^{-1}}) = l_1(\gamma_1) + l_1(\gamma_2) - 2l_1(\mathbf{p}).$$

Furthermore, since the isomorphism preserves the marked length spectrum, we have the relations:

- $l_2(\overline{\Phi(\gamma_i)}) = l_1(\gamma_i)$
- $l_2(\overline{\Phi(\gamma_2) * \Phi(\gamma_1^{-1})}) = l_1(\overline{\gamma_2 * \gamma_1^{-1}})$

Let $\overline{\Phi(\gamma_i)} = \eta_i$, so in particular, by Corollary 2.3, we have that $\Phi(\gamma_1) = \alpha\eta_1\alpha^{-1}$ and $\Phi(\gamma_2) = \beta\eta_2\beta^{-1}$ where α, β are geodesic paths in X_2 .

Claim 1. Using the notation from the previous paragraph, we must have: $\alpha = \beta$.

Indeed, if $\alpha \neq \beta$, we would have the inequalities:

$$l_2(\eta_1) + l_2(\eta_2) < l_2(\overline{\Phi(\gamma_2) * \Phi(\gamma_1^{-1})}) = l_1(\overline{\gamma_2 * \gamma_1^{-1}}) < l_1(\gamma_1) + l_1(\gamma_2).$$

But we have by the marked length spectrum being preserved, and the definition of η_i , that $l_2(\eta_i) = l_2(\overline{\Phi(\gamma_i)}) = l_1(\gamma_i)$ which gives us a contradiction (see Figure 1 for an illustration of this phenomena). Denote by $q \in X_2$ the endpoint of the path α . We can, without loss of generality, assume that $p_2 = q$ (by taking a change of basepoint for $\pi_1(X_2)$ if necessary). So we have reduced to the image being a pair of geodesic loops η_1 and η_2 based at q .

Claim 2. The geodesic loops η_1 and η_2 intersect in a path $\Phi\mathbf{p}$ of length precisely $l_1(\mathbf{p})$ passing through the point q (i.e. $\eta_1([0, l_1(\mathbf{p})]) = \eta_2([0, l_1(\mathbf{p})])$, but no such relationship holds for any larger interval).

In order to see this, let us assume that we can write $\eta_i = \sigma_i * \nu$, where ν is a path corresponding to the largest interval $[0, r]$ satisfying $\eta_1([0, r]) = \eta_2([0, r])$, and σ_i is the path $\eta_i([r, l_1(\gamma_i)])$ (in other words, ν is the longest path along which the two images curves agree, and σ_i is the rest of the respective curves). We claim that $l_2(\nu) = l_1(\mathbf{p})$.

By our choice of γ_1, γ_2 being $\mathbf{p}$ -distinguishing, we have the relation:

$$2l_1(\mathbf{p}) = l_1(\gamma_1) + l_1(\gamma_2) - l_1(\overline{\gamma_2 * \gamma_1^{-1}}) \quad (1)$$


Since we have that η_i are the geodesic loops in the free homotopy class of $\Phi(\gamma_i)$, and as our isomorphism preserves lengths, we have that:

$$l_2(\eta_i) = l_1(\gamma_i). \quad (2)$$

Furthermore, the composite $\gamma_2 * \gamma_1^{-1}$ corresponds to the composite $\eta_2 * \eta_1^{-1}$, which forces the equality $l_1(\overline{\gamma_2 * \gamma_1^{-1}}) = l_2(\overline{\eta_2 * \eta_1^{-1}})$, and the latter is freely homotopic to the geodesic loop $\sigma_2 * \sigma_1^{-1}$. This gives us that:

$$l_1(\overline{\gamma_1 * \gamma_2^{-1}}) = l_2(\overline{\eta_1 * \eta_2^{-1}}) = l_2(\eta_1) + l_2(\eta_2) - 2l_2(\nu). \quad (3)$$

Substituting equations (2) and (3) into equation (1), we obtain $2l_1(\mathbf{p}) = 2l_2(\nu)$ which immediately gives us the desired equality.

Figure 2: Two pairs of $\mathbf{p}$ -distinguishing curves.

Claim 3. The map is well-defined on geodesic paths (i.e. distinct pairs of distinguishing loops yield the same image path).

We have two possibilities, one of which is immediate: let $\gamma_1, \gamma_2, \gamma_3$ be geodesics based at p_1 an endpoint of $\mathbf{p}$ which are pairwise $\mathbf{p}$ -distinguishing. Then the three image geodesic loops $\overline{\Phi\gamma_1}, \overline{\Phi\gamma_2}, \overline{\Phi\gamma_3}$ are all based at q , and pairwise have the property that $\overline{\Phi\gamma_i}|_{[0, l_1(\mathbf{p})]} = \overline{\Phi\gamma_j}|_{[0, l_1(\mathbf{p})]}$. It is now immediate that they must all three coincide.

The other possibility to account for occurs if we have two distinct pairs (γ_1, γ_2) and (η_1, η_2) of $\mathbf{p}$ -distinguishing geodesics, but none of the pair (γ_i, η_j) are $\mathbf{p}$ -distinguishing. Since all four geodesics pass through $\mathbf{p}$, this means that the intersections $\gamma_i \cap \eta_j$ are all geodesic segments which extend the original $\mathbf{p}$ (see Figure 2 for an illustration of two such pairs near the geodesic). This immediately forces the geodesic loops to have a local picture near $\mathbf{p}$ as in Figure 2, with the pair (γ_1, γ_2) (locally) corresponding to the bottom left diagram, and the pair (η_1, η_2) (locally) corresponding to the bottom right diagram. Now focusing on the images of the four pairs of paths (γ_i, η_j) , we see that each pair gives image paths that coincide on a path of length precisely $l_1(\mathbf{p})$. Furthermore, the four image paths extend the common intersection by providing two branches on either side of the common intersection. In particular, the image paths give the exact same (top of Figure 2) local picture in X_2 . It is then easy to see that the corresponding pairs (γ_1, γ_2) and (η_1, η_2) both yield the common intersection, resulting in our map being well-defined. Now that we know that our map is well-defined, we can denote by $\Phi\mathbf{p}$ the image of the geodesic path $\mathbf{p}$ given by our construction.

*Claim 4. Let $\mathbf{p}_1$ and $\mathbf{p}_2$ be a pair of geodesically incident geodesic paths (by which we mean that the concatenated path $\mathbf{p}_1 * \mathbf{p}_2$ is also a geodesic path), meeting at a common vertex q which we will take as the basepoint for $\pi_1(X)$. Then the corresponding geodesic paths $\Phi\mathbf{p}_1$ and $\Phi\mathbf{p}_2$ are also geodesically incident.*




Figure 3: Incidence relations are preserved: the three possible cases.

Without loss of generality (by well-definedness of our map), we can assume that one of the closed loops used to find $\Phi \mathbf{p}_1, \Phi \mathbf{p}_2$ passes through both paths, hence can be used as one of both pairs. We refer to Figure 3 (top left) to illustrate our situation. In the top left figure, we have a pair of geodesically incident paths, with the big geodesic representing the common loop, and the two smaller ones intersecting the large one in $\mathbf{p}_1$ and $\mathbf{p}_2$ respectively.

Let us label (clockwise) the large common loop by γ , the two smaller ones by γ_1 and γ_2 respectively, and let each of the loops be oriented clockwise. Now consider the image loops (see Figure 3, remaining three pictures). If the resulting curves are not incident, we have that the two geodesic loops $\overline{\Phi(\gamma_1)}$ and $\overline{\Phi(\gamma_2)}$, which must intersect $\overline{\Phi(\gamma)}$, have intersections which do not represent incident subpaths of the geodesic loop $\overline{\Phi(\gamma)}$. We have three possible cases.

Firstly, the intersections with γ might be entirely disjoint (as in Figure 3, top right). Let $d > 0$ denote the distance between the two intersections, and note that we have:

$$l_2(\overline{\Phi(\gamma_1) * \Phi(\gamma_2)}) = l_2(\overline{\Phi(\gamma_1)}) + l_2(\overline{\Phi(\gamma_2)}) + 2d$$

But observe that the the geodesic loop $\overline{\gamma_1 * \gamma_2}$ has length which is bounded above by $l_1(\gamma_1) + l_1(\gamma_2)$, which combined with the fact that the isomorphism preserves the marked length spectrum, gives us a contradiction.

Another possibility is that the intersections might overlap in a subinterval of γ of

length d (as in Figure 3, bottom left). Well in that case, we can consider the geodesic loop corresponding to the composite curve $\gamma_1 * \gamma * \gamma_2$, and observe that it has length:

$$l_1(\overline{\gamma_1 * \gamma * \gamma_2}) = l_1(\gamma_1) + l_1(\gamma_2) + l_1(\gamma) - 2l_1(\mathbf{p}_1) - 2l_1(\mathbf{p}_2). \quad (4)$$

Looking at the geodesic loop corresponding to $\Phi(\gamma_1) * \Phi(\gamma) * \Phi(\gamma_2)$ in the image, we find that it has length:

$$l_2(\overline{\Phi(\gamma_1) * \Phi(\gamma) * \Phi(\gamma_2)}) = l_2(\Phi(\gamma_1)) + l_2(\Phi(\gamma_2)) + l_2(\Phi(\gamma)) - l_2(\Phi\mathbf{p}_1) - l_2(\Phi\mathbf{p}_2) + 2d.$$

And again, using that the isomorphism preserves the marked length spectrum, and comparing with equation (4) we get a contradiction.

Finally the third possibility is the one where one of the intersections lies entirely within the other intersection (Figure 3, bottom right). Without loss of generality, we can assume that $\overline{\Phi(\gamma_2)}$ is the small inner loop, while $\overline{\Phi(\gamma_1)}$ is the outer loop (so in particular $\Phi\mathbf{p}_2$ is a subpath of $\Phi\mathbf{p}_1$). If we let $0 < d < l_2(\Phi\mathbf{p}_1)$ be the distance between the left endpoints of $\Phi\mathbf{p}_i$, a simple calculation will show that:

$$l_2(\overline{\Phi(\gamma_1) * \Phi(\gamma) * \Phi(\gamma_2)}) = l_2(\overline{\Phi(\gamma_1)}) + l_2(\overline{\Phi(\gamma)}) + l_2(\overline{\Phi(\gamma_2)}) - 2d - 2l_2(\Phi\mathbf{p}_2)$$

which we can compare with the expression in equation (4). Using the fact that the isomorphism preserves lengths, this immediately yields that $d = l_1(\mathbf{p}_1)$, which again is a contradiction.

This gives us that the image paths $\Phi\mathbf{p}_1$ and $\Phi\mathbf{p}_2$ are subpaths of the geodesic loop γ which agree at one endpoint, but not in any larger neighborhood of the endpoint. Since γ is cyclically reduced, this immediately forces the concatenation $\Phi\mathbf{p}_1 * \Phi\mathbf{p}_2$ to be a geodesic path, hence geodesically incident paths map to geodesically incident paths. Observe that it is now clear that if $\mathbf{p} = \mathbf{q}_2 * \mathbf{q}_1$ is a geodesic path, written as a concatenation of subpaths, then $\Phi\mathbf{p} = \Phi\mathbf{q}_2 * \Phi\mathbf{q}_1$ (since the $\mathbf{q}_i$ are geodesically incident).

Claim 5. Let $\mathbf{p}_1$ and $\mathbf{p}_2$ be a pair of incident geodesic paths, meeting at a common vertex q which we will take as the basepoint for $\pi_1(X)$. Then the corresponding geodesic paths $\Phi\mathbf{p}_1$ and $\Phi\mathbf{p}_2$ are also incident.

To see this, note that if the incident paths are *geodesically* incident, we are done by the previous claim. So let us assume not. Then by Lemma 2.1, we have that the reduced path corresponding to the concatenation $\mathbf{p}_2 * \mathbf{p}_1$ is of the form $\mathbf{r}_2 * \mathbf{r}_1$, where $\mathbf{r}_i$ is a subpath of $\mathbf{p}_i$. Furthermore, $\mathbf{p}_2 = \mathbf{r}_2 * \mathbf{q}^{-1}$, while $\mathbf{p}_1 = \mathbf{q} * \mathbf{r}_1$. In particular, from the previous claim, we have that $\Phi\mathbf{p}_2 = \Phi\mathbf{r}_2 * \Phi\mathbf{q}^{-1}$ and $\Phi\mathbf{p}_1 = \Phi\mathbf{q} * \Phi\mathbf{r}_1$. However, reversal of paths is preserved under the map Φ we have constructed (since

we can take the same pair of distinguishing loops with reversed orientation). This immediately yields our last claim.

Observe that, since incidence relations are preserved, our map induces a well defined bijection between the set of endpoints, as any two paths joining a pair of branch points can be viewed as incident paths (at both ends), hence will map to incident paths (again at both ends). Finally, we note that for any pair of branch points in X_1 we can find a geodesic joining the points which realizes the distance between them (namely, the distance minimizer, by Corollary 2.3). As a result, we have that the distance between the image points is precisely the same. Hence we have an isometry. This completes the proof of Proposition 3.1

Finally, we provide a proof of our main theorem:

Proof: (Theorem 3.1). By Proposition 3.1, we already know that there is an isometry between the sets of branch points. So all we need to show is that we can extend this isometry to a global isometry. First we note that, since we are working with complete spaces, an isometry between subsets extends to an isometry between their closures. So we are left with extending our map to points that do not lie in the closure of the branch points.

So let $p \in \text{Conv}(X_1)$ lie outside of the closure of the branching locus. Then there exists a metric neighborhood of p with no branch points. Now let U_p be a largest neighborhood (maximal under containment) of p which does not contain any branch points. Then U_p must be isometric to an interval of some length r . Furthermore, the boundary of this interval in Y_1 consists solely of branch points, and this interval defines a path between them. By the proof of Proposition 3.1, there is a corresponding pair of branch points in X_2 , and a path of exactly the same length between them. Furthermore, the corresponding path in X_2 cannot have any branch points either (else applying the construction from Proposition 3.1 to the inverse would yield a branch point in our original path). So we can extend our map by prescribing it to be an isometry from each U_p to the corresponding (branchless) path in X_2 . It is now immediate to verify that this does indeed give an isometry.

Finally, using completeness, we can extend once more to get an isometric map from Y_1 into X_2 . Note that, by construction, this map will in fact have image lying in Y_2 , hence our map is an isometric embedding from Y_1 into Y_2 . But note that we can apply the same construction to ϕ^{-1} , yielding an isometric embedding from Y_2 into Y_1 . Furthermore, the composite of the two maps corresponds to the map from Y_1 to itself obtained by applying this construction to the identity isomorphism. Hence, must be the identity map on Y_1 . This implies the map is an isometry from Y_1 to Y_2 . Finally, by a result of Curtis & Fort (Corollary 2.1 in [7]), we have that $\pi_1(Y_i)$ injects in $\pi_1(X_i)$, and by the naturality of the construction, we see that the map we

constructed induces the isomorphism (up to change of basepoint) between the $\pi_1(Y_i)$. The theorem follows.

Finally, we use our main theorem to get a somewhat stronger result:

Proposition 3.2 (Generalization of Main Theorem). *Let X_1, X_2 be a pair of complete, compact, one dimensional geodesic spaces. If they have the same marked length spectrum, then there is an isometry from $\overline{\text{Conv}}(X_1)$ to $\overline{\text{Conv}}(X_2)$*

Proof. Let us denote by $Y_i \subseteq X_i$ the π_1 -hull of X_i . We want to prove that Y_1 is isometric to Y_2 . Without loss of generality, we can assume that the basepoints for the fundamental groups $\pi_1(X_i)$ actually lie in the Y_i . Now a result of Curtis & Fort (Corollary 2.1 in [7]) implies that $\pi_1(Y_i)$ is actually a subgroup of $\pi_1(X_i)$.

4 Group actions on $\mathbb{R}$ -trees.

For completeness, we indicate how our theorem overlaps with the results of Culler & Morgan [6] (a more general version of their work is contained in the paper by Alperin & Bass [1]). We begin by reviewing some terminology for group actions on $\mathbb{R}$ -trees.

Definition 4.1. A group action on an $\mathbb{R}$ -tree is said to be *minimal* if there are no proper invariant subtrees.

Definition 4.2. Let G be a group acting by isometries on an $\mathbb{R}$ -tree. The action is said to be *semi-simple* if either of the following three conditions hold:

- there is a G -invariant line in T
- every element of G fixes a point
- every element of G acts fixed point freely, there is no G -invariant line, and no G -invariant end of T .

In this case, we call the action type I, II, or III respectively.

Definition 4.3. The translation length function associated to an action of G on an $\mathbb{R}$ -tree T is the function on G defined by $l(g) := \inf_{x \in T} d(x, g(x))$.

The main result of Culler-Morgan [6] then reads:

Theorem 4.1. *Let $T_1 \times G \longrightarrow T_1$ and $T_2 \times G \longrightarrow T_2$ be a pair of minimal semi-simple actions of a group G on $\mathbb{R}$ -trees T_1 and T_2 with the same translation length function. Then there exists an equivariant isometry from T_1 to T_2 .*

We can derive this result for type III semi-simple actions from Theorem 3.1, with the added hypothesis that the action is orientation preserving and acts properly discontinuously. The argument is the following:

Proof: (Weak form of Theorem 4.1). We start by recalling that, for an $\mathbb{R}$ -tree T , any isometry g either has a fixed point or has positive translation length. Furthermore, if g is fixed point free, then there is a unique line in T that is g -invariant. Assume now that we have a pair of minimal type III semi-simple actions of a group G on a pair of $\mathbb{R}$ -trees with the same translation length function. We claim that there exists an equivariant isometry from T_1 to T_2 .

Consider the quotient spaces $X_i = T_i/G$ with induced metric. We start by noting that the covering dimension cannot increase under quotients, and since the quotients are connected, they both have $Cdim = 1$. Furthermore, as the action is properly discontinuous, $\pi_1(X_i) = G$, and the translation length function on G corresponds precisely to the length function of loops in the quotient space X_i .

We now have to show that the spaces X_i are π_1 -convex (i.e. using the notation of our theorem, $Y_i = X_i$). If not, we have a subset $Y \subset X_i$ with the property that every geodesic loop in X_i actually lies in Y . Now consider the lift $\tilde{Y} \subset T_i$ to the $\mathbb{R}$ -tree. We claim that $\tilde{Y}$ is connected. Indeed, Y is connected, so considering a single connected component Y_i of $\tilde{Y}$, we can lift paths to see that Y_i contains pre-images of *every* point in Y . Now assume that there are two distinct connected components Y_1, Y_2 of $\tilde{Y}$. Observe that there is an element $g \in G$ that takes Y_1 to Y_2 . Consider a geodesic loop γ in Y representing g , and observe that every point in γ (by definition of π_1 -convex) must lie in Y . Now let $p \in \gamma$, and consider a pair of points p_1, p_2 in Y_1, Y_2 mapping down to the same point $p \in Y$. Lifting γ to T_i , we obtain a path joining p_1 to p_2 which consists purely of points in the pre-image of Y . Hence Y_1 and Y_2 are in fact in the same path connected component of $\tilde{Y}$. We conclude that $\tilde{Y}$ consists of a single path connected component.

Next we note that a path connected subset of an $\mathbb{R}$ -tree is itself an $\mathbb{R}$ -tree. Hence $\tilde{Y}$ is a proper subtree of T . Furthermore, since $\tilde{Y}$ consists of all pre-images of points in Y , this is a G -invariant subtree. By minimality of the G -action, this forces $\tilde{Y} = T_i$. But this in turn forces $Y = X_i$. This concludes our proof.

Finally, Theorem 3.1 implies that X_1 is in fact isometric to X_2 . Lifting this isometry, we obtain a G -equivariant isometry between the respective trees T_1 and T_2 .

The reason why we need proper discontinuity of the action is precisely because, in the absence of proper discontinuity, we have no way to relate the fundamental group of the quotient to the original group. Conversely, if we have a pair of geodesic length spaces of topological dimension one which are both semi-locally simply connected, then our result follows from theirs (by considering the fundamental group acting on the universal cover). However, there are numerous examples of geodesic length spaces

of topological dimension one which are not semi-locally simply connected (Hawaiian Earring, Menger curve, Sierpinski curve, etc), for which our result does not *a priori* follow from theirs.

5 Bibliography

- [1] Alperin, R. & Bass, H. *Length functions of group actions on Λ -trees*. in Combinatorial group theory and topology (Alta, Utah, 1984), pp. 265–378, Ann. of Math. Stud., 111, Princeton Univ. Press, Princeton, NJ, 1987.
- [2] Bridson, M. R. & Haefliger, A. *Metric spaces of non-positive curvature*. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 319. Springer-Verlag, Berlin, 1999. xxii+643 pp.
- [3] Burago, D., Burago, Y. & Ivanov, S. *A course in metric geometry*. Graduate Studies in Mathematics, 33. American Mathematical Society, Providence, RI, 2001. xiv+415 pp.
- [4] Cannon, J. & Conner, G. *On the fundamental group of one-dimensional spaces*. (Preprint), <http://www.math.byu.edu/conner/preprints/index.html>.
- [5] Croke, C. B. *Rigidity for surfaces of nonpositive curvature*. Comment. Math. Helv. 65 (1990), no. 1, pp. 150–169.
- [6] Culler, M. & Morgan, J. W. *Group actions on $\mathbf{R}$ -trees*. Proc. London Math. Soc. (3) 55 (1987), no. 3, pp. 571–604.
- [7] Curtis, M. L. & Fort, M. K., Jr. *The fundamental group of one-dimensional spaces*. Proc. Amer. Math. Soc. 10 (1959), pp. 140–148.
- [8] Curtis, M. L. & Fort, M. K., Jr. *Homotopy groups of one-dimensional spaces*. Proc. Amer. Math. Soc. 8 (1957), pp. 577–579.
- [9] Dal’Bo, F. & Kim, I. *Marked length rigidity for symmetric spaces*. Comment. Math. Helv. 77 (2002), no. 2, pp. 399–407.
- [10] Hamenstädt, U. *Entropy-rigidity of locally symmetric spaces of negative curvature*. Ann. of Math. (2) 131 (1990), no. 1, pp. 35–51.
- [11] Hurewicz, W. & Wallman, H. *Dimension Theory*. Princeton Mathematical Series, v. 4. Princeton University Press, Princeton, N. J., 1941. vii+165 pp.
- [12] Otal, J.-P. *Le spectre marqué des longueurs des surfaces à courbure négative*. (French) [The marked spectrum of the lengths of surfaces with negative curvature] Ann. of Math. (2) 131 (1990), no. 1, pp. 151–162.